

Electronic Supplementary Material to: Dynamical Predictability of Leading Interannual Variability Modes of the Asian-Australian Monsoon in Climate Models*

Lin WANG¹, Hong-Li REN^{1,2}, Fang ZHOU³, Nick DUNSTONE⁴, and Xiangde XU¹

¹State Key Laboratory of Severe Weather, and Institute of Tibetan Plateau Meteorology,
Chinese Academy of Meteorological Sciences, Beijing 100081, China

²Department of Atmospheric Science, School of Environmental Studies,
China University of Geosciences, Wuhan 430074, China

³Collaborative Innovation Center on Forecast and Evaluation of Meteorological Disasters/Key Laboratory of
Meteorological Disaster, Ministry of Education/Joint International Research Laboratory of Climate and Environment
Change, Nanjing University of Information Science and Technology, Nanjing 210044, China

⁴Met Office Hadley Centre, FitzRoy Road, Exeter, Devon, EX1 3PB, United Kingdom

ESM to: Wang, L., H.-L. Ren, F. Zhou, N. Dunstone, and X. D. Xu, 2023: Dynamical predictability of leading interannual variability modes of the Asian-Australian Monsoon in climate models. *Adv. Atmos. Sci.*, **40**(11), 1998–2012, <https://doi.org/10.1007/s00376-023-2288-2>.




Fig. S1. The first SEOF spatial patterns of AAM U850 anomalies (units: m s^{-1}) as derived from different initial seasons in BCC_CSM1.1M data: (a) SEOF1-frMAM; (b) SEOF1-frJJA; (c) SEOF1-frSON; (d) SEOF1-frDJF.




Fig. S2. Same as Fig. S1, but for the second SEOF.